

**Total RNA Purification Micro Kit**  
Product # 35300

**Product Insert**

Norgen's Total RNA Purification Micro Kit provides a rapid and sensitive method for the isolation and purification of total RNA from small input amounts of cultured animal cells, tissue samples, and microdissected samples including laser-capture microdissection (LCM). The kit purifies all sizes of RNA, from large mRNA and ribosomal RNA down to microRNA (miRNA) and small interfering RNA (siRNA). The RNA is preferentially purified from other cellular components such as proteins without the use of phenol or chloroform. The purified RNA is of the highest integrity, and can be used in a number of downstream applications including real time PCR, reverse transcription PCR, Northern blotting, RNase protection and primer extension, and expression array assays.

**Norgen's Purification Technology**

Purification is based on spin column chromatography using Norgen's proprietary resin as the separation matrix. The RNA is preferentially purified from other cellular components such as proteins without the use of phenol or chloroform. The process involves first lysing the cells or tissue of interest with the provided Buffer RL (please see the flow chart on page 4). Ethanol is then added to the lysate, and the solution is loaded onto a micro spin-column. Norgen's resin binds RNA in a manner that depends on ionic concentrations. Thus only the RNA will bind to the column, while the contaminating proteins will be removed in the flowthrough or retained on the top of the resin. The bound RNA is then washed with the provided Wash Solution A in order to remove any remaining impurities, and the purified total RNA is eluted with the Elution Solution A. The special design of the micro spin-column allows a small elution volume of as little as 20 µL. The purified RNA is of the highest integrity, and can be used in a number of downstream applications.

**Specifications**

Kit Specifications	
Maximum Column Binding Capacity	35 µg
Maximum Column Loading Volume	650 µL
Minimum Elution Volume	20 µL
Size of RNA Purified	All sizes, including small RNA (<200 nt)
Maximum Amount of Starting Material:	
Animal Cells	5 x 10 <sup>5</sup> cells
Animal Tissues	3 mg (for most tissues*)
Laser-Captured Microdissection (LCM)	Up to 5 x 10 <sup>5</sup> cells
Time to Complete 10 Purifications	20 minutes
Average Yields	
HeLa Cells (1 x 10 <sup>5</sup> cells)	1.5 µg

\* for fibrous tissue, an additional Proteinase K treatment is required

### Advantages

- Fast and easy processing using rapid micro spin-column format
- Small elution volume of 20  $\mu$ L
- Isolate total RNA, from large rRNA down to microRNA (miRNA) without compromising total yield
- No phenol or chloroform extractions
- Isolate high quality total RNA from a variety of sources
- RNA can be isolated and detected from as little as a single animal cell

### Kit Components

Component	Product # 35300 (50 preps)
Buffer RL	40 mL
Wash Solution A	38 mL
Elution Solution A	6 mL
Micro Spin Columns	50
Collection Tubes	50
Elution tubes (1.7 mL)	50
Product Insert	1

### Storage Conditions and Product Stability

All solutions should be kept tightly sealed and stored at room temperature. These reagents should remain stable for at least 1 year in their unopened containers.

### Precautions and Disclaimers

This kit is designed for research purposes only. It is not intended for human or diagnostic use.

Ensure that a suitable lab coat, disposable gloves and protective goggles are worn when working with chemicals. For more information, please consult the appropriate Material Safety Data Sheets (MSDSs). These are available as convenient PDF files online at [www.norgenbiotek.com](http://www.norgenbiotek.com).

The **Buffer RL** contains guanidine salts, and should be handled with care. Guanidine salt forms highly reactive compounds when combined with bleach, thus care must be taken to properly dispose of any of these solutions.

## Customer-Supplied Reagents and Equipment

You must have the following in order to use the Total RNA Purification Kit:

### *For All Protocols*

- Benchtop microcentrifuge
- 96 - 100% ethanol
- $\beta$ -mercaptoethanol (optional)

### *For Animal Cell Protocol*

- PBS (RNase-free)

### *For Animal Tissue Protocol*

- Liquid nitrogen
- Mortar and pestle
- 70% ethanol

### *For Laser-Captured Microdissection (LCM) Protocol*

- Sterile fine forceps
- Water bath or heat block set at 42°C
- 70% ethanol

## Working with RNA

RNases are very stable and robust enzymes that degrade RNA. Autoclaving solutions and glassware is not always sufficient to actively remove these enzymes. The first step when preparing to work with RNA is to create an RNase-free environment. The following precautions are recommended as your best defense against these enzymes.

- The RNA area should be located away from microbiological work stations
- Clean, disposable gloves should be worn at all times when handling reagents, samples, pipettes, disposable tubes, etc. It is recommended that gloves are changed frequently to avoid contamination
- There should be designated solutions, tips, tubes, lab coats, pipettes, etc. for RNA only
- All RNA solutions should be prepared using at least 0.05% DEPC-treated autoclaved water or molecular biology grade nuclease-free water
- Clean all surfaces with commercially available RNase decontamination solutions
- When working with purified RNA samples, ensure that they remain on ice during downstream applications

## Flowchart

Procedure for Purifying Total RNA using Norgen's Total RNA Purification Micro Kit

Lyse cells or tissue using **Buffer RL**



Add Ethanol



Bind to column

**SPIN**



Wash three times  
with Wash Solution A

**SPIN**



Elute RNA with  
Elution Solution A

**SPIN**



**Purified Total RNA**

## Procedures

All centrifugation steps are carried out in a benchtop microcentrifuge. Various speeds are required for different steps, so please check your microcentrifuge specifications to ensure that it is capable of the proper speeds. All centrifugation steps are performed at room temperature. The correct rpm can be calculated using the formula:

$$RPM = \sqrt{\frac{RCF}{(1.118 \times 10^{-5}) (r)}}$$

where  $RCF$  = required gravitational acceleration (relative centrifugal force in units of  $g$ );  $r$  = radius of the rotor in cm; and  $RPM$  = the number of revolutions per minute required to achieve the necessary  $g$ -force.

## Section 1. Preparation of Lysate From Various Cell Types

### Notes Prior to Use

- The steps for preparing the lysate are different depending on the starting material (**Step 1**). However, the subsequent steps are the same in all cases (**Steps 2 – 6**).
- Please ensure that the correct procedure for preparing the lysate from your starting material is followed.
- All centrifugation steps are carried out in a benchtop microcentrifuge at 14,000 x  $g$  (~ 14,000 RPM) except where noted. All centrifugation steps are performed at room temperature.
- A variable speed centrifuge should be used for maximum kit performance. If a variable speed centrifuge is not available a fixed speed centrifuge can be used, however reduced yields may be observed.
- Ensure that all solutions are at room temperature prior to use.
- Prepare a working concentration of the **Wash Solution A** by adding 90 mL of 96 - 100% ethanol (provided by the user) to the supplied bottle containing the concentrated **Wash Solution A**. This will give a final volume of 128 mL. The label on the bottle has a box that may be checked to indicate that the ethanol has been added.
- **Optional:** The use of  $\beta$ -mercaptoethanol in lysis is highly recommended for most animal tissues, particularly those known to have a high RNase content (ex: pancreas), including LCM samples. It is also recommended for users who wish to isolate RNA for sensitive downstream applications. Add 10  $\mu$ L of  $\beta$ -mercaptoethanol (provided by the user) to each 1 mL of Buffer RL required.  $\beta$ -mercaptoethanol is toxic and should be dispensed in a fume hood. Alternatively, the Buffer RL can be used as provided.
- It is important to work quickly during this procedure.

## 1A. Lysate Preparation from Cultured Animal Cells

### Notes Prior to Use

- The maximum recommended input of cells is  $5 \times 10^5$ . A hemocytometer can be used in conjunction with a microscope to count the number of cells. As a general guideline, each well of a confluent 12-well plate of HeLa cells will contain  $5 \times 10^5$  cells.
- Cell pellets can be stored at  $-70^\circ\text{C}$  for later use or used directly in the procedure. Determine the number of cells present before freezing.
- Frozen pellets should be stored for no longer than 2 weeks to ensure that the integrity of the RNA is not compromised.
- Frozen cell pellets should not be thawed prior to beginning the protocol. Add the Buffer RL directly to the frozen cell pellet (**Step 1A(ii) c**).

### 1A(i). Cell Lysate Preparation from Cells Growing in a Monolayer

- a. Aspirate media and wash cell monolayer with an appropriate amount of PBS. Aspirate PBS.
- b. Add 350  $\mu$ L of **Buffer RL** directly to culture plate.
- c. Lyse cells by gently tapping culture dish and swirling buffer around plate surface for five minutes.
- d. Transfer lysate to a microcentrifuge tube.
- e. Add 200  $\mu$ L of 96 – 100% ethanol (provided by the user) to the lysate. Mix by vortexing for 10 seconds. **Proceed to Step 2.**

### 1A (ii). Cell Lysate Preparation from Cells Growing in Suspension and Lifted Cells

- a. Transfer cell suspension to an RNase-free tube (not provided) and centrifuge at no more than 200  $\times g$  (~2,000 RPM) for 10 minutes to pellet cells.
- b. Carefully decant the supernatant.

**Note:** For inputs of over  $10^5$  cells, 5-10  $\mu$ L of media may be left behind with the pellet in order to ensure that the pellet is not dislodged. For inputs of fewer than  $10^5$  cells, 30-50  $\mu$ L of media may be left behind in order to ensure that the pellet, which could be invisible, is not dislodged.

- c. Add 350  $\mu$ L of **Buffer RL** to the pellet. Lyse cells by vortexing for 15 seconds. Ensure that the entire pellet is completely dissolved before proceeding to the next step.
- d. Add 200  $\mu$ L of 96 - 100% ethanol (provided by the user) to the lysate. Mix by vortexing for 10 seconds. **Proceed to Step 2.**

## 1B. Lysate Preparation from Animal Tissues

### Notes Prior to Use

- Norgen's Total RNA Purification Micro Kit is designed for isolating RNA from small amounts of non-fibrous tissue samples (up to 3 mg in most cases). If a larger amount of starting material or fibrous tissue is desired, an additional Proteinase K treatment is required. Please refer to Appendix A for instruction. Also, to isolate total RNA from larger amounts of tissue Norgen also offers a Total RNA Purification Kit (Cat# 17200) and an Animal Tissue RNA Purification Kit (Cat# 25700).
- RNA in animal tissues is not protected after harvesting until it is disrupted and homogenized. Thus it is important that the procedure is carried out as quickly as possible, particularly the Cell Lysate Preparation step.
- Fresh or frozen tissues may be used for the procedure. Tissues should be flash-frozen in liquid nitrogen and transferred immediately to a  $-70^{\circ}\text{C}$  freezer for long-term storage. Tissues may be stored at  $-70^{\circ}\text{C}$  for several months. When isolating total RNA from frozen tissues ensure that the tissue does not thaw during weighing or prior to grinding with the mortar and pestle.
- Tissues stored in RNA stabilization reagents such as *RNAlater*<sup>®</sup> are compatible with this isolation procedure. Prior to isolation, carefully remove the tissue from the storage reagent using forceps, and dry excessive liquid.
- The maximum recommended input of tissue varies depending on the type of tissue being used. Please refer to Table 1 below as a guideline for maximum tissue input amounts. If your tissue of interest is not included in the table below we recommend starting with an input of no more than 3 mg.

**Table 1. Recommended Maximum Input Amounts of Different Tissues**

Tissue	Maximum Input Amount
Brain, Kidney, Liver, Lung, Spleen	3 mg
Heart, Muscle	Refer to Appendix A

**1B. Cell Lysate Preparation from Animal Tissues**

- a. Excise the tissue sample from the animal.
- b. Determine the amount of tissue by weighing. Please refer to Table 1 for the recommended maximum input amounts of different tissues. For tissues not included in the table, we recommend starting with an input of no more than 3 mg.
- c. Transfer the tissue into a mortar that contains an appropriate amount of liquid nitrogen to cover the sample. Grind the tissue thoroughly using a pestle.
- d. Allow the liquid nitrogen to evaporate, without allowing the tissue to thaw.
- e. Add 400  $\mu$ L of **Buffer RL** to the tissue sample and continue to grind until the sample has been homogenized. Homogenize by passing the lysate 5-10 times through a 25 gauge needle attached to a syringe.
- f. Using a pipette, transfer the lysate into an RNase-free microcentrifuge tube (not provided).
- g. Spin the lysate for 2 minutes to pellet any cell debris. Transfer the supernatant to another RNase-free microcentrifuge tube. Note the volume of the supernatant/lysate.
- h. Add an equal volume of 70% ethanol (provided by the user) to the lysate volume collected (100  $\mu$ L of ethanol is added to every 100  $\mu$ L of lysate). Vortex to mix. **Proceed to Step 2.**

**1C. Lysate Preparation from Laser-Captured Microdissection (LCM)**

**Notes Prior to Use**

- LCM samples obtained from frozen sections are recommended. Formalin-Fixed, Paraffin-Embedded sections may also be used. However, RNA isolated from FFPE samples generally has poorer quality than that from frozen sections.

**1C. Cell Lysate Preparation from Laser-Captured Microdissection (LCM)**

- a. Aliquot 300  $\mu$ L of **Buffer RL** to an RNase-free microcentrifuge tube.
- b. Remove the thermoplastic film containing the captured cells using sterile fine forceps. Carefully submerge the sample into the aliquoted **Buffer RL**. Close the microcentrifuge cap.
- c. Incubate the sample at 42°C for 30 minutes. Apply vortex for 15 seconds after every 10 minutes.
- d. At the end of the incubation, vortex the tube one more time for 15 seconds. The thermoplastic film may be removed at this point using sterile fine forceps. Otherwise, proceed to **Step 1Ce**.
- e. Add 300  $\mu$ L of 70% ethanol (provided by the user) to the lysate. Vortex to mix. **Proceed to Step 2.**

## Section 2. Total RNA Purification from All Types of Lysate

**Note:** The remaining steps of the procedure for the purification of total RNA are the same from this point forward for all the different types of lysate.

### 2. Binding RNA to Column

- a. Assemble a micro spin-column with one of the provided collection tubes
- b. Apply up to 600  $\mu\text{L}$  of the lysate with the ethanol (from **Step 1**) onto the column and centrifuge for 1 minute at  $\geq 3,500 \times g$  (~6,000 RPM).

**Note:** Ensure the entire lysate volume has passed through into the collection tube by inspecting the column. If the entire lysate volume has not passed, spin for an additional minute at  $14,000 \times g$  (~14,000 RPM).

- c. Discard the flowthrough. Reassemble the spin column with its collection tube.
- d. Depending on your lysate volume, repeat Step **2b** and **2c** as necessary.

### Optional Step:

Norgen's Total RNA Purification Mirco Kit isolates total RNA with minimal amounts of genomic DNA contamination. However, an optional **On-Column DNA Removal Protocol** is provided in Appendix B for maximum removal of residual DNA that may affect sensitive downstream applications. It is recommended that Norgen's RNase-Free DNase I Kit (Product # 25710) be used for this step. This step should be performed at this point in the protocol.

### 3. Column Wash

- a. Apply 400  $\mu\text{L}$  of **Wash Solution A** to the column and centrifuge for 1 minute.

**Note:** Ensure the entire Wash Solution A has passed through into the collection tube by inspecting the column. If the entire wash volume has not passed, spin for an additional minute.

- b. Discard the flowthrough and reassemble the spin column with its collection tube.
- c. Repeat steps **3a** and **3b** to wash column a second time.
- d. Wash column a third time by adding another 400  $\mu\text{L}$  of **Wash Solution A** and centrifuging for 1 minute.
- e. Discard the flowthrough and reassemble the spin column with its collection tube.
- f. Spin the column for 2 minutes in order to thoroughly dry the resin. Discard the collection tube.

### 4. RNA Elution

- a. Place the column into a fresh 1.7 mL Elution tube provided with the kit.
- b. Add 40  $\mu\text{L}$  of **Elution Solution A** to the column.

**Note:** For higher concentrations of RNA, a lower elution volume may be used. A minimum volume of 20  $\mu\text{L}$  is recommended

- c. Centrifuge for 2 minutes at **200 x g (~2,000 RPM)**, followed by 1 minute at **14,000 x g (~14,000 RPM)** Note the volume eluted from the column. If the entire volume has not been eluted, spin the column at  $14,000 \times g$  (~14,000 RPM) for 1 additional minute.

**Note:** For maximum RNA recovery, it is recommended that a second elution be performed into a separate microcentrifuge tube (Repeat **Steps 4b and 4c**).



## 5. Storage of RNA

The purified RNA sample may be stored at  $-20^{\circ}\text{C}$  for a few days. It is recommended that samples be placed at  $-70^{\circ}\text{C}$  for long term storage.

## Appendix A

### Lysate Preparation from Animal Tissues with the use of Proteinase K

#### *Customer-Supplied Reagent*

- RNase-Free Proteinase K
- RNase-Free Water
- $\beta$ -mercaptoethanol

#### *Notes Prior to Use*

- Ensure that all solutions are at room temperature prior to use.
- We recommend the use of Norgen's **Proteinase K** (Cat # 17904) for this step. Reconstitute each of the **Proteinase K** vials in 600  $\mu\text{L}$  of molecular biology grade water or 10 mM Tris.HCl pH 7.5 (RNase-Free). For every isolation, 20  $\mu\text{L}$  of the reconstituted Proteinase K is needed. Aliquot the remainder into small fractions and store the unused portions at  $-20^{\circ}\text{C}$  until needed.
- If using another source of **Proteinase K**, reconstitute in molecular biology grade water or 10 mM Tris.HCl pH 7.5 (RNase-Free) to give a 20 mg/mL final concentration. For every isolation, 20  $\mu\text{L}$  of the reconstituted Proteinase K is needed. Aliquot the remainder into small fractions and store the unused portions at  $-20^{\circ}\text{C}$  until needed.
- Add 10  $\mu\text{L}$  of  $\beta$ -mercaptoethanol (provided by the user) to each 1 mL of **Buffer RL** required.  $\beta$ -mercaptoethanol is toxic and should be dispensed in a fume hood.
- RNA in animal tissues is not protected after harvesting until it is disrupted and homogenized. Thus it is important that the procedure is carried out as quickly as possible, particularly the Cell Lysate Preparation step.
- Fresh or frozen tissues may be used for the procedure. Tissues should be flash-frozen in liquid nitrogen and transferred immediately to a  $-70^{\circ}\text{C}$  freezer for long-term storage. Tissues may be stored at  $-70^{\circ}\text{C}$  for several months. When isolating total RNA from frozen tissues ensure that the tissue does not thaw during weighing or prior to grinding with the mortar and pestle.
- Tissues stored in RNA stabilization reagents such as RNA<sup>later</sup><sup>®</sup> are compatible with this isolation procedure. Prior to isolation, carefully remove the tissue from the storage reagent using forceps, and dry any excessive liquid.
- This protocol is particularly suitable for isolating RNA from up to 7.5 mg of tissues including fibrous, connective tissues.

#### **Cell Lysate Preparation**

- a. Excise the tissue sample from the animal.
- b. Determine the amount of tissue by weighing.
- c. Transfer the tissue into a mortar that contains an appropriate amount of liquid nitrogen to cover the sample. Grind the tissue thoroughly using a pestle.

**Note:** The use of liquid nitrogen is recommended. However, if homogenization without flash-freezing is preferred, proceed to Step **e**.

- d. Allow the liquid nitrogen to evaporate, without allowing the tissue to thaw.
- e. Add 300  $\mu\text{L}$  of **Buffer RL** to the tissue sample and continue to grind until the sample has been homogenized.

**Note:** Maximum homogenization may be achieved by passing the lysate 5-10 times through a 25 gauge needle attached to a syringe.

- f. Using a pipette, transfer the lysate into an RNase-free microcentrifuge tube (not provided).
- g. Add 600  $\mu\text{L}$  of **RNase-Free Water** (not provided) to the lysate. Vortex to mix.
- h. Add 20  $\mu\text{L}$  of reconstituted Proteinase K to the lysate, and incubate at 55°C for 15 minutes. Vortex the tubes occasionally during incubation.
- i. Spin the lysate for 2 minutes to pellet any cell debris. Transfer the supernatant to another RNase-free microcentrifuge tube.
- j. Add 450  $\mu\text{L}$  of 96 - 100% ethanol (provided by the user) to the lysate. Vortex to mix.  
**Proceed to Step 2.**

## Appendix B

### Protocol for Optional On-Column DNA Removal

Norgen's Total RNA Purification Micro Kit isolates total RNA with minimal amounts of genomic DNA contamination. However, an optional protocol is provided below for maximum removal of residual DNA that may affect sensitive downstream applications. It is recommended that Norgen's RNase-Free DNase I Kit (Product # 25710) be used for this step.

1. For every on-column reaction to be performed, prepare a mix of 15  $\mu\text{L}$  of **DNase I** and 100  $\mu\text{L}$  of **Enzyme Incubation Buffer** using Norgen's RNase-Free DNase I Kit (Product # 25710). Mix gently by inverting the tube a few times. **DO NOT VORTEX.**

**Note:** If using an alternative DNase I, prepare a working stock of 0.25 Kunitz unit/ $\mu\text{L}$  RNase-free DNase I solution according to the manufacturer's instructions. A 100  $\mu\text{L}$  aliquot is required for each column to be treated.

2. Perform the appropriate Total RNA Isolation Procedure for your starting material up to and including "**Binding to Column**" (Steps 1 and 2 of all protocols)
3. Apply 400  $\mu\text{L}$  of **Wash Solution A** to the micro spin column and centrifuge for 2 minute. Discard the flowthrough. Reassemble the spin column with its collection tube.
4. Apply 100  $\mu\text{L}$  of the RNase-free DNase I solution prepared in Step 1 to the column and centrifuge at 14, 000 x g (~14 000 RPM) for 1 minute.

**Note:** Ensure that the entire DNase I solution passes through the column. If needed, spin at 14, 000 x g (~14 000 RPM) for an additional minute.

5. After the centrifugation in Step 4, pipette the flowthrough that is present in the collection tube back onto the top of the column.

**Note:** Ensure Step 5 is performed in order to ensure maximum DNase activity and to obtain maximum yields of RNA, in particular for small RNA species.

6. Incubate the column assembly at 25 - 30°C for 15 minutes.
7. Without any further centrifugation, proceed directly to the second wash step in the "**Column Wash**" section (Step 3c).

## Troubleshooting Guide

Problem	Possible Cause	Solution and Explanation
Poor RNA Recovery	Incomplete lysis of cells or tissue	Ensure that the appropriate amount of Buffer RL was used for the amount of cells or tissue.
	Column has become clogged	Do not exceed the recommended amounts of starting materials. The amount of starting material may need to be decreased if the column shows clogging below the recommended levels. See also “Clogged Column” below.
	An alternative elution solution was used	It is recommended that the Elution Solution A supplied with this kit be used for maximum RNA recovery.
	Ethanol was not added to the lysate	Ensure that the appropriate amount of ethanol is added to the lysate before binding to the column.
	Ethanol was not added to the Wash Solution A	Ensure that 90 mL of 96 - 100% ethanol is added to the supplied Wash Solution A prior to use.
	Low RNA content in cells or tissues used	Different tissues and cells have different RNA contents, and thus the expected yield of RNA will vary greatly from these different sources. Please check literature to determine the expected RNA content of your starting material.
	Cell Culture: Cell monolayer was not washed with PBS LCM: Sample was not incubated at 42°C for 30 minutes	Ensure that the cell monolayer is washed with the appropriate amount of PBS in order to remove residual media from cells. Ensure that the incubation at 42°C for the removal and lysis of cells from the thermoplastic film.
Clogged Column	Insufficient solubilization of cells or tissues	Ensure that the appropriate amount of lysis buffer was used for the amount of cells or tissue.
	Maximum number of cells or amount of tissue exceeds kit specifications	Refer to specifications to determine if amount of starting material falls within kit specifications
	High amounts of genomic DNA present in sample	The lysate may be passed through a 25 gauge needle attached to a syringe 5-10 times in order to shear the genomic DNA prior to loading onto the column.

<b>Problem</b>	<b>Possible Cause</b>	<b>Solution and Explanation</b>
Clogged Column	Centrifuge temperature too low	Ensure that the centrifuge remains at room temperature throughout the procedure. Temperatures below 15°C may cause precipitates to form that can cause the columns to clog.
RNA is Degraded	RNase contamination	RNases may be introduced during the use of the kit. Ensure proper procedures are followed when working with RNA. Please refer to “ <i>Working with RNA</i> ” at the beginning of this user guide.
	Procedure not performed quickly enough	In order to maintain the integrity of the RNA, it is important that the procedure be performed quickly. This is especially important for the Cell Lysate Preparation Step in the Animal Tissue protocol, since the RNA in animal tissues is not protected after harvesting until it is disrupted and homogenized.
	Improper storage of the purified RNA	For short term storage RNA samples may be stored at –20°C for a few days. It is recommended that samples be stored at –70°C for longer term storage.
	Frozen tissues or cell pellets were allowed to thaw prior to RNA isolation	Do not allow frozen tissues to thaw prior to grinding with the mortar and pestle in order to ensure that the integrity of the RNA is not compromised.
	Starting material may have a high RNase content	For starting materials with high RNAase content, it is recommended that β-mercaptoethanol be added to the Buffer RL.
RNA does not perform well in downstream applications	RNA was not washed 3 times with the provided Wash Solution A	Traces of salt from the binding step may remain in the sample if the column is not washed 3 times with Wash Solution A. Salt may interfere with downstream applications, and thus must be washed from the column.
	Ethanol carryover	Ensure that the dry spin under the Column Wash procedure is performed, in order to remove traces of ethanol prior to elution. Ethanol is known to interfere with many downstream applications.
Genomic DNA contamination	Large amounts of starting material used	Perform RNase-free DNaseI digestion on the RNA sample after elution to remove genomic DNA contamination. It is recommended that Norgen’s RNase-Free DNase I Kit (Product # 25710) be used for this step.

Related Products	Product #
Proteinase K – 2 Vials	17904
RNase-Free DNase I Kit	25710
Total RNA Purification Kit	17200
Total RNA Purification 96-Well Kit	24300
Total RNA Purification Maxi Kit	26800
Animal Tissue RNA Purification Kit	25700
Plant/Fungi Total RNA Purification Kit	25800
RNA/Protein Purification Kit	24100
RNA/DNA/Protein Purification Kit	24000
Cytoplasmic & Nuclear RNA Purification Kit	21000
Leukocyte RNA Purification Kit	21200
microRNA Purification Kit	21300
100b RNA Ladder	15002
1kb RNA Ladder	15003

### Technical Support

Contact our Technical Support Team between the hours of 8:30 and 5:30 (Eastern Standard Time) at (905) 227-8848 or Toll Free at 1-866-667-4362.

Technical support can also be obtained from our website ([www.norgenbiotek.com](http://www.norgenbiotek.com)) or through email at [techsupport@norgenbiotek.com](mailto:techsupport@norgenbiotek.com).

3430 Schmon Parkway, Thorold, ON Canada L2V 4Y6  
 Phone: (905) 227-8848  
 Fax: (905) 227-1061  
 Toll Free in North America: 1-866-667-4362