

TREATS UP TO
5,000 SQ FT*

GUARANTEED RESULTS†

Spectracide®

GRUB STOP® Once & Done! + Fertilizer

LARVACIDA MÁS
FERTILIZANTE

KILLS GRUBS NOW
Prevents Them All Season Long!

- ▶ Helps Your Lawn Stay Strong During Stressful Summer Months
- ▶ Also Kills European Crane Fly Larvae and Mole Crickets
- ▶ Apply Anytime Spring Through Late Summer



ESPAÑOL
VEA EL REVERSO

14-0-3

ACTIVE INGREDIENT:
Imidacloprid (CAS# 138261-41-3)0.22% KEEP OUT OF REACH OF CHILDREN MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
INERT INGREDIENTS99.78% CAUTION See back for additional precautionary statements. PRECAUCIÓN Consulte la parte posterior para conocer las precauciones adicionales.

NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASSAU, SUFFOLK, KINGS AND QUEENS COUNTIES, NEW YORK.

*Treats up to 5,000 sq ft for grubs
†For terms of guarantee, see back panel.

Net Wt 14.35 lb (6.5 kg)

Apply with
Spreader

Spectracide®

GRUB STOP® Once & Done! + Fertilizer

TREATS UP TO
5,000 SQ FT*

LARVACIDA MÁS
FERTILIZANTE



GRUB STOP[®]

Once & Done!

+ Fertilizer LARVACIDA MÁS FERTILIZANTE

TREATS UP TO
5,000 SQ FT*

Lifecycle of the Grub



MARCH - MAY

In spring, grubs awake from hibernation to feed on grass roots, destroying your lawn.



JUNE - JULY

In summer, beetles emerge to feed on your garden... and lay their eggs.



AUGUST - OCTOBER

By fall, eggs hatch into grubs, which starts the destruction cycle over again... by feeding on your lawn's roots.

Product Facts

What It Does:

Kills Grubs, European Crane Fly Larvae and Mole Crickets.

Where To Use:

For use on Lawns, Ground Covers, Flower Beds, and around Trees & Shrubs.

When To Use:

Apply anytime spring through late summer.

Questions & Comments:

Call 1-800-917-5438 or visit our website at www.spectracide.com



STOP READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not apply near fish pools, ponds, streams or lakes. Do not use in vegetable gardens.

GENERAL INFORMATION

Spectracide[®] Grub Stop[®] Once & Done!™ plus Fertilizer is for use on lawns, flower beds, ground covers, trees and shrubs to control white grubs (including larvae of Asiatic garden beetle, black turfgrass ataenius, chafers, Japanese beetle, May or June beetle, and Oriental beetle), mole crickets and European crane fly larvae.

For best results, apply this product before grubs, mole crickets or European crane fly larvae hatch and can cause any damage. Contact your County Extension Agent for the best dates to apply in your area.

Pests Controlled	When to Apply	Indication of Damage
Grubs	Apply once anytime from May through mid-August.	White grubs cause widespread damage to lawns by feeding on grass roots, causing it to turn yellow, then brown and die.
European crane fly larvae	Apply once when these insects are present in large numbers, late summer to mid-fall.	European crane fly larvae cause damage by feeding on grass roots and disturbing the soil surface.
Mole crickets	Apply once anytime from mid-May through mid-June. Florida residents – apply once anytime between early April through early May.	Mole crickets cause widespread damage by tunneling through the soil, while some species also feed on roots. Tunneling through the soil uproots the grass causing it to dry out, become thin and weedy, turn yellow and eventually die.

LAWNS

Application Instructions

See Spreader Settings chart. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace, making parallel passes. Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Water thoroughly after application.

For best results:

- Apply early before grubs, mole crickets and European crane fly larvae hatch.
- Irrigate thoroughly within 24 hours of application to form a protective zone in the soil.
- Avoid mowing until granules are watered into the soil.
- Do not apply to water-saturated areas. This may decrease product effectiveness.
- This product can be applied over soil or mulch. Remove black plastic or other weed barrier material prior to application.

Keep children and pets off treated area until it has dried.

FLOWER BEDS, GROUND COVERS, TREES AND SHRUBS

Application Instructions

Apply this product at the rate of 2 Tbsp per 10 sq ft of planting bed (1/4 cups per 100 sq ft of planting bed). Evenly spread granules to the soil over flower beds, ground cover plantings, shrub beds and under trees. The use of a handheld spreader is recommended. Rake granules into the soil around plants. Water thoroughly after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

Pesticide Disposal: Do not reuse empty bag. Place empty bag in trash.

IMPORTANT: It is best to use the entire product in accordance with label directions. Never pour insecticide down a drain or storm sewer. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

Have product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

If swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

In case of a medical emergency call Poison Control Center at 1-800-917-5438.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

*Treats up to 5,000 sq ft for grubs.

GUARANTEED ANALYSIS

Total Nitrogen (N) 14.0%

14% Urea Nitrogen*

Soluble Potash (K₂O) 3.0%

Derived from: Urea and Muriate of Potash.

F1074

MADE IN THE USA Distributed by Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642
EPA Reg. No. 53883-204-845
EPA Est. Nos. 9198-AL-1(A), 9198-OH-2(B), 9198-OH-1(M),
44616-MO-1(C), 44616-GA-1(D), 572-NJ-1(T), 8669-AL-1(B),
8660-OH-1(B), 9688-MO-1(I), 39578-TX-1(A), 69208-MO-1(A)
Circled letter is first letter of lot number. 11-10868 © 2007 UIC

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond United Industries Corporation's control it is impossible for United Industries to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED INDUSTRIES MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of United Industries is authorized to make any warranty beyond those contained herein or to modify the warranties contained herein. United Industries disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT UNITED INDUSTRIES' ELECTION, THE REPLACEMENT OF PRODUCT.

No contamine el agua ni los alimentos para personas o animales por medio del almacenamiento o la eliminación.

Almacenamiento del pesticida: Guarde en su envase original en un lugar fresco y seco, que no sea accesible a los niños, de preferencia en un gabinete cerrado bajo llave. No permita que se congele.

Eliminación del pesticida: No vuelva a utilizar la bolsa vacía. Tire la bolsa vacía en la basura.

IMPORTANTE: Es mejor usar todo el producto de acuerdo con las instrucciones de la etiqueta. Nunca vierta un insecticida por un desague ni por una alcantarilla pluvial. Si es necesario desechar el producto sin usar, por favor siga todas las normas locales y estatales pertinentes.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua ni los alimentos para personas o animales por medio del almacenamiento o la eliminación.

Almacenamiento del pesticida: Guarde en su envase original en un lugar fresco y seco, que no sea accesible a los niños, de preferencia en un gabinete cerrado bajo llave. No permita que se congele.

Eliminación del pesticida: No vuelva a utilizar la bolsa vacía. Tire la bolsa vacía en la basura.

IMPORTANTE: Es mejor usar todo el producto de acuerdo con las instrucciones de la etiqueta. Nunca vierta un insecticida por un desague ni por una alcantarilla pluvial. Si es necesario desechar el producto sin usar, por favor siga todas las normas locales y estatales pertinentes.

PRECAUCIONES

RIESGOS PARA LOS SERES HUMANOS Y LOS ANIMALES DOMÉSTICOS

PRECAUCIÓN: Nocivo si se ingiere o se absorbe por la piel. Provoca irritación ocular. Evite el contacto con la piel, los ojos o la ropa.

Después de manipular el producto, lávese muy bien con agua y jabón.

PRIMEROS AUXILIOS Cuando llame a un centro de control de envenenamientos o a un médico, intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.

Si se ingiere: • Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

• Si la persona puede tragar, haga que beba poco a poco un vaso de agua.

• No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico.

• No administre nada por la boca a una persona que haya perdido el conocimiento.

Si entra en contacto con la piel o la ropa: • Quite la ropa contaminada.

• Lave inmediatamente la piel con abundante agua durante 15-20 minutos.

• Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Si entra en contacto con los ojos: • Mantenga los ojos abiertos y enjuague con agua ligeramente, con cuidado, durante 15 a 20 minutos.

• Si hay lentes de contacto, retírelos después de 5 minutos, después continúe enjuagando los ojos.

• Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

En caso de una urgencia médica, llame al centro de control de envenenamientos al 1-800-917-5438.

RIESGOS PARA EL MEDIO AMBIENTE

Este pesticida es altamente tóxico para los invertebrados acuáticos. No aplique directamente al agua. No contamine el agua al lavar el equipo o cuando elimine las aguas de lavado del equipo.



GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address on this bag, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

PROOF OF PURCHASE

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-sb.com>.

FPO UPC
0 71121 95718 4

TREATS UP TO
5,000 SQ
FT*

GUARANTEED RESULTS[†]

Spectracide

GRUB STOP®

Once & Done!

+Fertilizer

LARVACIDA MÁS
FERTILIZANTE

KILLS GRUBS NOW
Prevents Them All Season Long!

- ▶ Helps Your Lawn Stay Strong During Stressful Summer Months
- ▶ Also Kills European Crane Fly Larvae and Mole Crickets
- ▶ Apply Anytime Spring Through Late Summer

ESPAÑOL
VEA EL REVERSO

14-0-3

ACTIVE INGREDIENT:

Imidacloprid (CAS# 138261-41-3)0.22%

KEEP OUT OF REACH OF CHILDREN

INERT INGREDIENTS 99.78%

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

TOTAL: 100.00%

CAUTION See back for additional

PRECAUCIÓN Consulte la parte posterior para

conocer las precauciones adicionales.

NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASSAU, SUFFOLK, KINGS AND QUEENS COUNTIES, NEW YORK.

*Treats up to 5,000 sq ft for grubs

[†]For terms of guarantee, see back panel.

Net Wt 14.35 lb (6.5 kg)



Apply with
Spreader

Spectracide

GRUB STOP[®]
Once & Done!
+ Fertilizer

LARVACIDA MÁS
FERTILIZANTE

TREATS UP TO
5,000 SQ FT*

Spectracide

GRUB STOP®

Once & Done!™

+ Fertilizer

TREATS UP TO
5,000 SQ FT*

Lifecycle of the Grub



MARCH - MAY

In spring, grubs awake from hibernation to feed on grass roots, destroying your lawn.



Grubs change into pupae in late spring.



JUNE - JULY
In summer, beetles emerge to feed on your garden...



...and lay their eggs.



AUGUST - OCTOBER
By fall, eggs hatch into grubs, which starts the destruction cycle over again...
...by feeding on your lawn's roots.

Product Facts

What It Does:

Kills Grubs, European Crane Fly Larvae and Mole Crickets.

Where To Use:

For use on Lawns, Ground Covers, Flower Beds, and around Trees & Shrubs.



When To Use:

Apply anytime spring through late summer.

Questions & Comments:

Call 1-800-917-5438 or visit our website at www.spectracide.com



STOP READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not apply near fish pools, ponds, streams or lakes. Do not use in vegetable gardens.

GENERAL INFORMATION

Spectracide® Grub Stop® Once & Done!™ plus Fertilizer is for use on lawns, flower beds, ground covers, trees and shrubs to control white grubs (including larvae of Asiatic garden beetle, black turfgrass ataenius, chafers, Japanese beetle, May or June beetle, and Oriental beetle), mole crickets and European crane fly larvae.

For best results, apply this product before grubs, mole crickets or European crane fly larvae hatch and can cause any damage. Contact your County Extension Agent for the best dates to apply in your area.

Pests Controlled	When to Apply	Indication of Damage
Grubs	Apply once anytime from May through mid-August.	White grubs cause widespread damage to lawns by feeding on grass roots, causing it to turn yellow, then brown and die.
European crane fly larvae	Apply once when these insects are present in large numbers, late summer to mid-fall.	European crane fly larvae cause damage by feeding on grass roots and disturbing the soil surface.
Mole crickets	Apply once anytime from mid-May through mid-June. Florida residents – apply once anytime between early April through early May.	Mole crickets cause widespread damage by tunneling through the soil, while some species also feed on roots. Tunneling through the soil uproots the grass causing it to dry out, become thin and weedy, turn yellow and eventually die.

LAWNS

Application Instructions

See Spreader Settings chart. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace, making parallel passes. Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Water thoroughly after application.

For best results:

- Apply early before grubs, mole crickets and European crane fly larvae hatch.
- Irrigate thoroughly within 24 hours of application to form a protective zone in the soil.

SPREADER SETTINGS FOR LAWN APPLICATION AJUSTES DEL DIFUSOR PARA LA APLICACIÓN EN CÉSPED

Rate for grubs and European crane fly larvae: 2.87 lb per 1,000 sq ft
Rate for mole crickets: 3.8 lb per 1,000 sq ft



Broadcast/Rotary Spreaders 2.87 lb/1,000 sq ft 3.8 lb/1,000 sq ft

Scotts® Broadcast/Rotary Spreaders

Scotts® Basic™, Deluxe™, EdgeGuard™

CÉSPEDES

Instrucciones de aplicación

Vea el cuadro de Ajustes del difusor. Regule el difusor en el ajuste apropiado. Asegúrese de que el difusor funcione correctamente y distribuya las gránulos uniformemente. Aplique al borde y después llene en el centro para asegurarse de que toda el área quede completamente cubierta. Para asegurarse de que el producto se distribuya uniformemente, camine a ritmo regular, pasando el producto en forma paralela. No use un difusor que aplique los gránulos en franjas estrechas, concentradas o irregulares. Riegue abundantemente después de la aplicación.

Para obtener mejores resultados:

- Aplique antes de que los gusanos, los grillos cebolleros y las larvas de las tipulas europeas salgan.
- Riegue bien durante las 24 horas posteriores a la aplicación para formar una zona de protección en la tierra.

that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace, making parallel passes. Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Water thoroughly after application.

For best results:

- Apply early before grubs, mole crickets and European crane fly larvae hatch.
- Irrigate thoroughly within 24 hours of application to form a protective zone in the soil.
- Avoid mowing until granules are watered into the soil.
- Do not apply to water-saturated areas. This may decrease product effectiveness.
- This product can be applied over soil or mulch. Remove black plastic or other weed barrier material prior to application.

Keep children and pets off treated area until it has dried.

FLOWER BEDS, GROUND COVERS, TREES AND SHRUBS

Application Instructions

Apply this product at the rate of 2 Tbsp per 10 sq ft of planting bed (1/4 cups per 100 sq ft of planting bed). Evenly spread granules to the soil over flower beds, ground cover plantings, shrub beds and under trees. The use of a handheld spreader is recommended. Rake granules into the soil around plants. Water thoroughly after application.

SPREADER SETTINGS FOR LAWN APPLICATION AJUSTES DEL DIFUSOR PARA LA APLICACIÓN EN CÉSPED



Rate for grubs and European crane fly larvae: 2.87 lb per 1,000 sq ft

Rate for mole crickets: 3.8 lb per 1,000 sq ft

Broadcast/Rotary Spreaders	2.87 lb/1,000 sq ft	3.8 lb/1,000 sq ft
Scotts® Broadcast/Rotary Spreaders		
Scotts Basic™, Deluxe™ EdgeGuard®	6 1/2	7 1/2
Lawn Pro™, SpeedyGreen® or Standard™	11 1/2	12
Earthway EV-N-SPRED® 2050P		
Drop Spreaders		
Scotts Drop Spreaders	7 1/2	8 1/4
Scotts AccuGreen®		

Note: 1 Tbsp = 0.28 oz; 1 cup = 4.43 oz

Typically a lawn utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied at 2.87 lb per 1,000 sq ft, this product provides 0.4 lb of nitrogen per 1,000 sq ft; and when applied at the 3.8 lb per 1,000 sq ft rate, this product provides 0.53 lb of nitrogen per 1,000 sq ft. Contact your local Extension Service for information specific to your geographic region.

En general, un césped utiliza entre 1 y 4 lb. de nitrógeno cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica a la dosis de 2.87 lb. cada 1,000 pies cuadrados, este producto proporciona 0.4 lb. de nitrógeno cada 1,000 pies cuadrados; y cuando se aplica a la dosis de 3.8 lb. cada 1,000 pies cuadrados, este producto proporciona 0.53 lb. de nitrógeno cada 1,000 pies cuadrados. Comuníquese con la oficina local del servicio de extensión agrícola para obtener información específica para la región donde vive.

diffusor funcione correctamente y distribuya los gránulos uniformemente. Aplique al borde y después llene en el centro para asegurarse de que toda el área quede completamente cubierta. Para asegurarse de que el producto se distribuya uniformemente, camine a ritmo regular, pasando el producto en forma paralela. No use un difusor que aplique los gránulos en franjas estrechas, concentradas o irregulares. Riegue abundantemente después de la aplicación.

Para obtener mejores resultados:

- Aplique antes de que los gusanos, los grillos cebolleros y las larvas de las típulas europeas salgan.
- Riegue bien durante las 24 horas posteriores a la aplicación para formar una zona de protección en la tierra.
- Evite cortar hasta que los gránulos se diluyan en la tierra.
- No aplique en áreas saturadas de agua. Esto podría disminuir la eficacia del producto.
- Este producto se puede aplicar sobre tierra o mantillo. Antes de la aplicación retire el plástico negro u otra barrera para las malezas.

Mantenga a los niños y a las mascotas fuera de las áreas tratadas hasta que el producto se haya secado.

LECHOS DE FLORES, CUBRESUELOS, ÁRBOLES Y ARBUSTOS

Instrucciones de aplicación

Aplique este producto a la dosis de 2 cucharadas soperas cada 10 pies cuadrados de lecho de plantación (1/4 taza por cada 100 pies cuadrados de lecho de plantación). Distribuya los gránulos de manera uniforme en la tierra sobre lechos de flores, cubresuelos, lechos de arbustos y debajo de árboles. Se recomienda el uso de un difusor manual. Introduzca los gránulos en la tierra, alrededor de las plantas, con el rastrillo. Riegue abundantemente después de la aplicación.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

Pesticide Disposal: Do not reuse empty bag. Place empty bag in trash.

IMPORTANT: It is best to use the entire product in accordance with label directions. Never pour insecticide down a drain or storm sewer. If it is necessary to dispose of unused product, please follow any applicable state and local guidelines.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

Have product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

If swallowed:

- Call a Poison Control Center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a Poison Control Center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin with plenty of water for 15-20 minutes.
- Call a Poison Control Center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

In case of a medical emergency call Poison Control Center at 1-800-917-5438.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

*Treats up to 5,000 sq ft for grubs.

GUARANTEED ANALYSIS 14-0-3

Total Nitrogen (N) 14.0%
14% Urea Nitrogen*

Soluble Potash (K₂O) 3.0%

Derived from: Urea and Muriate of Potash.

F1074



Distributed by Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 53883-204-8845

EPA Est. Nos. 9198-AL-1(A), 9198-OH-2(B), 9198-OH-1(M),
44616-MO-1(C), 44616-GA-1(D), 572-NJ-1(T), 8660-AL-1(BN),
8660-OH-1(BD), 9688-MO-1(U), 39578-TX-1(AM), 69208-MO-1(AO)

Circled letter is first letter of lot number.

11-10868

© 2007 UIC

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond United Industries Corporation's control it is impossible for United Industries to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED INDUSTRIES MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of United Industries is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. United Industries disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT UNITED INDUSTRIES' ELECTION, THE REPLACEMENT OF PRODUCT.

PRIMEROS AUXILIOS

Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.

Si se ingiere:

- Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
- Si la persona puede tragar, haga que beba poco a poco un vaso de agua.
- No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico.
- No administre nada por la boca a una persona que haya perdido el conocimiento.

Si entra en contacto con la piel o la ropa:

- Quite la ropa contaminada.
- Lave inmediatamente la piel con abundante agua durante 15-20 minutos.
- Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Si entra en contacto con los ojos:

- Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos.
- Si hay lentes de contacto, retírelos después de los primeros 5 minutos, después continúe enjuagando los ojos.
- Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

En caso de una urgencia médica, llame el centro de control de envenenamientos al 1-800-917-5438.

RIESGOS PARA EL MEDIO AMBIENTE

Este pesticida es altamente tóxico para los invertebrados acuáticos. No aplique directamente al agua. No contamine el agua al lavar el equipo o cuando elimine las aguas de lavado del equipo.



GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address on this bag, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

PROOF OF PURCHASE

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-sb.com>.

FPO UPC
0 71121 95718 4