Net Weight 12.5 LB | Peso Neto 5.6 kg

COVERS UP TO 5,000 SQ. FT.

Boosts Nutrient Uptake for a lush, green Lawn

Increases Drought Tolerance & Reduces Watering Needs by Extending Root Absorption Area

KILLS OVER 250 WEEDS DOWN TO THE ROOT

35-0-4

WEED & FEED II

PENNINGTON®

Pennington®

ENHANCED FERTILIZER

Feed More Lawn with Less Fertilizer

Color Blocks

Top Seal Customer 15mm

CUT HERE TO OPEN

Cut / Tear Here

The Pennington® UltraGreen® Difference

Half of the nitrogen in this premium lime™ is stabilized™, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow release for up to 3 months of extended feeding. Additionally, these unique fertilizer blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients. These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

Pennington® UltraGreen® Weed & Feed, 35-0-4 WILL NOT BURN when used as directed. We're so confident you'll love the results of Pennington® UltraGreen® Weed & Feed, 30-0-4 that it comes with the Pennington Satisfaction Guarantee.†

†excluding Pennington® UltraGreen® Starter Fertilizer

Benefits of Pennington® UltraGreen® Weed & Feed, 30-0-4

- Kills over 250 weeds including Henbit, Dollar Weed, White Clover, Plantain and Chickweed
- Feeds for up to 3 months
- Rich in nitrogen for lush, green grass
- Enriched with potassium for overall plant vigor
- Feeds more than 5,000 sq. ft.
- Ready-to-Use Fertilizer

STOP! READ THE ENTIRE LABEL FIRST!

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for Treatment. You may also contact 1-800-383-0926 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use on dichondra, carpetgrass, 'Floratam' St. Augustinegrass, creeping bentgrass, or turfgrass when dichondra clovers are present.
- Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
- Do not use this product on or near decorative plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy.
- Apply only by actively growing turf. Check with your local Cooperative Extension Agency to obtain specific information on local turf bed management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

INSTRUCTIONS:

Pennington® UltraGreen® Weed & Feed, 30-0-4 is a combination of three proven weed killers – 2,4-D, mecoprop, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit and sprange. In addition, a 30-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS:

- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, patio, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Reseeding or over seeding areas may occur three (3) weeks after the application of this product.

HOW MUCH TO APPLY: For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass; centipedegrass, hybrid bermudagrass, and colonial bentgrass: Apply 2.5 pounds of product per 1,000 square feet of turfgrass. If needed, a second application may be made after 30 days or more. The maximum number of broadcast applications per treatment site is 2 per year. Do not apply more than 5 pounds of product per 1,000 sq. ft. per year. Typically a lawn utilizes 3 to 4 pounds of nitrogen per 1,000 sq. ft. per growing season. When applied as directed, this product provides 0.75 pounds of nitrogen per 1,000 sq. ft. Contact your local Extension Service for information specific to your geographic region.

WHEN TO APPLY: Apply when broadleaf weeds are actively growing in spring or early fall. Apply when the soil is cold (choose a cool day) to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS:

1. Mow lawn to remove top third (not to exceed 1 by 2 days before application.
2. Water lawn thoroughly (at least 1 to 2 days before application to scatter moisture until the root watering (see Step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not water lawn weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTINGS: Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand. Coverage – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as non-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

Bag Size (lbs)	Application Rate (lbs / 1,000 sq. ft.)	Treated Area for this Bag Size (sq. ft.)
12.5	2.5	5,000

SINGLE PASS DISTRIBUTION:

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS: Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions. To provide proper distribution, calibrate spreader before application. For 2.5 pounds of product per 1,000 square feet of turfgrass. Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shifting off the spreader before each turn at edge of border strips.

SETTING YOUR LAWN SPREADER

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and wind speed can affect how fast the lawn food comes out. Watch flow rate. If fertilizer appears to be coming out either slower or faster than you are covering your lawn, adjust the setting. Close the spreader when filling, turning or stopping.

Spreader	Setting Delivers 2.5 lbs / 1000 sq. ft.
Scotts® Turf Builder® Classic Drop	3.5
Scotts® Turf Builder® Edgeguard® Mini	3.5
Scotts Broadcast	3.25
Fleming Deep® Broadcast	3

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

WARRANTY & DISCLAIMER: To the extent consistent with applicable law, Central Garden & Pet, Garden Division guarantees this fine product to the extent of the purchase price, only when used according to directions.

30-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	30%
10% Urea Nitrogen*	
Soluble Potash (K2O)	4%
Calcium (Ca)	1%

*Derived from Urea, Polymer Coated Urea, Mixtures of Potash, and Limestone.

*Contains 12% Slowly Available urea nitrogen from polymer coated urea

*Contains 18% nitrogen stabilized with N-(n-butyl) thiophosphoric triamide

**Stabilized nitrogen minimizes nitrogen losses to the environment and enhances its efficiency.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

SOIL AMENDING GUARANTEED ANALYSIS

Active Ingredients:

Mycorrhizal Fungi	1.12 pounds per
Glomus intraradices	0.28 pounds per
Glomus mosseae	0.28 pounds per
Glomus aculeatum	0.28 pounds per
Glomus etunicatum	0.28 pounds per

Inert Ingredients (as a soil amendment):

1% 100% Inert ingredients

1.184% Herbicide ingredients

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Certain states have limits on fertilizer application rates and timing. Consult your State Cooperative Extension for current information on annual application limitations for lawn fertilizers in your region.

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

NOTICE: This product contains chemicals known to the state of California to cause cancer and birth defects or other reproductive harm.

1 Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to:

Pennington Brand
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

La diferencia de Pennington® UltraGreen®

Los productos Pennington® UltraGreen® son más eficientes que otros fertilizantes. La mitad del nitrógeno de esta lime™ de calidad está estabilizado™, lo que permite que la hierba aproveche mejor los nutrientes que usted agregue y reduce la pérdida de nutrientes hacia el ambiente. La otra mitad del nitrógeno es de liberación lenta hasta por 3 meses de nutrición extendida. Por otra parte, estas exclusivas mezclas contienen hongos micorrizas, unos organismos naturales que han formado parte de los suelos vigorosos durante millones de años. Estos hongos microscópicos aumentan la superficie de absorción de las raíces, mejorando así grandemente la capacidad del césped para tomar agua y los recursos vitales del suelo como son el agua y los nutrientes. Estos pequeños gránulos de fertilizante están empacados con nutrientes esenciales para plantas y hongos, para que usted pueda nutrir más césped con menos fertilizante.

†excluye el fertilizante de hierba Pennington® UltraGreen® Starter Fertilizer

Ventajas del fertilizante y nutriente Pennington® UltraGreen® de 30-0-4

- Elimina más de 250 malezas incluyendo trébol blanco, amiño de lewis, trébol blanco, llantén y álcine
- Cinco para el tratamiento de los céspedes norteños y sureños*
- Revenerdimiento rápido y nutrición extendida
- No contiene fitotoxinas
- Hasta 3 meses de nutrición
- Rico en nitrógeno para lograr un césped verde y copioso
- Enriquecido con potasio para aportarle vigor general a las plantas

*No se debe usar en la hierba de San Agustín 'Floratam'

El fertilizante y nutriente Pennington® UltraGreen®, de 30-0-4 NO QUEMA la hierba cuando se usa como se indica. Confíamos tanto en que le encantarán los resultados del uso del fertilizante y nutriente Pennington® UltraGreen®, de 30-0-4 que le ofrecemos la garantía de satisfacción de Pennington.†

¡DETÉNGASE! LA PRIMERO LA ETIQUETA COMPLETA. PRESTE ATENCIÓN A TODAS LAS PRECAUCIONES Y CUMPLA DETENIDAMENTE CON TODAS LAS INDICACIONES.

PRECAUCIONES

Peligros para personas y animales domésticos

PRECAUCIÓN: Causa irritación moderada de los ojos. Evite el contacto con los ojos o la ropa. Use camisa de manga larga, pantalones largos, zapatos y medias. Lávese bien con agua y jabón después de manipular el producto y antes de comer, beber, masticar goma, consumir tabaco o ir al baño.

Primera ayuda

En caso de contacto con los ojos	• Mantenga el ojo abierto y enjuague lento y suavemente con aproximadamente 15-20 minutos. Si se le retiran de contacto, retírelos después de los primeros 5 minutos y continúe enjuagando ocularmente. • Llame a un centro de control de envenenamiento o un médico para recibir consejo sobre el tratamiento.

Tenga a mano el recipiente o la etiqueta del producto cuando llame al centro de control de envenenamiento o el médico, o cuando busca usted el tratamiento. Para recibir información sobre el tratamiento médico de emergencia, llame también al 1-800-383-0926.

Peligros ambientales

Este pesticida es tóxico para los peces y invertebrados acuáticos y puede afectar de manera adversa a las plantas para las que no está destinado. Para proteger el ambiente, no permita que el pesticida entre en las alcantarillas para las tormentas, zanjas de drenaje, canalletas o aguas de superficie. Aplique este producto en tiempo de calma, cuando no se pronostique lluvia en viento por las próximas 24 horas para garantizar que no vuelva o la lluvia no trasladen el pesticida fuera del área tratada. Si el producto cae en la entrada de carros, sendero, acera o calle, barrido hacia el césped o jardín para evitar que sea arrastrado hacia corrientes de agua o sistemas de drenaje. Este producto quimico tiene propiedades y características asociadas con sustancias quimicas detectadas en las aguas subterraneas. El uso de este producto en areas donde el suelo es permeable puede ocasionar la contaminación de las aguas subterraneas, especialmente si el nivel acuífero o hidrico no es profundo. La aplicación alrededor de cisternas o pozos puede ocasionar la contaminación del agua potable o el agua subterranea.

INSTRUCCIONES DE USO

El uso de este producto de manera diferente a lo indicado en su etiqueta es una violación de las leyes federales.

RESTRICCIONES DE USO:

- No aplique este producto de manera que entre en contacto con ninguna persona o mascota, ya sea directa o indirectamente. Mantenga a las personas y mascotas fuera del área durante la aplicación. No permita que las personas o mascotas entren en el área tratada hasta que el polvo se haya asentado.
- No se debe usar en céspedes cultivados para la venta o otro uso comercial como tepes, producción comercial de semillas o para fines de investigación.
- No lo use en céspedes de gramínea, zacate amargo, hierba de San Agustín 'Floratam', agrostide rastrera o dichondra cuando trébol de hoja hidrate desaparezcan.
- No aplique este producto a hierbas recién sembradas hasta que no estén bien asentadas o después de la segunda o la tercera poda.
- No utilice este producto en flores, vegetales, plantas de cobertura de terreno ni plantas de jardinería o decoración exteriores, incluyendo arbustos y árboles. No exceda la dosis especificada para cualquier área y no aplique este producto dentro de la línea de gotas ni sobre las raíces superficiales de árboles, arbustos y otras plantas ornamentales.
- No utilice este producto en o cerca de sus jardines ornamentales y evite la propagación de los gránulos hacia las raíces expuestas o brotes adventicios (rizomas) dentro de la línea de gotas de las raíces de los árboles y arbustos deseables, ya que puede causar daño.
- No lo aplique directamente o cerca del agua, después que caiga la tormenta, canaletas, alcantarillas o zanjas de drenaje. No lo aplique a una distancia menor de 25 pies de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías ni océanos. No aplique cuando haya viento.
- Aplique solo a céspedes en crecimiento. Verifique con la Agencia de Extensión Cooperativa de su comunidad para obtener información específica acerca de los regímenes prácticos del cuidado del césped. Verifique con el gobierno de su condado o ciudad para determinar si hay restricciones locales sobre el uso o la aplicación de fertilizantes.

INSTRUCCIONES:

El fertilizante y nutriente Pennington® UltraGreen®, de 30-0-4 es una combinación de tres herbicidas con efectividad demostrada: 2,4-D, Mecoprop-p y Dicamba. Este herbicida controla una amplia gama de malezas del césped, incluyendo diente de león, álcine, centinodia, llantén, artigo mayor y plantón. Además, el fertilizante de 30-0-4 provee un buen surtido de nutrientes para plantas.

PRECAUCIONES DE USO:

- Aplique este producto directamente a su césped y barra el producto que caiga en la entrada de carros, acera, canalleta o calle hacia el área tratada. Para evitar que el producto sea arrastrado, no riegue en exceso la área tratada de manera que el agua se salga del lugar y no aplique cuando lluvia o cuando se espere lluvia.
- No resembre o sobresembre de semilla se puede realizar a las tres (3) semanas después de la aplicación de este producto.

CUANDO APLICARLO: Para usar en pasto pará, festuca, bulbo perenne, hierba Bermuda común, hierba bahía, zoysia, hierba de San Agustín, campiés, híbridos de hierba Bermuda y nudo de dos. Aplique 2.5 lbs del producto por 1000 pies cuadrados de césped. De ser necesario, se puede hacer una segunda aplicación después de 30 días o más. El número máximo de aplicaciones por dispersión para cada área de tratamiento es de 2 por año. No aplique más de 5 libras del producto por cada 1000 pies cuadrados por año. Generalmente, un césped consume de 3 a 4 libras de nitrógeno por cada 1000 pies cuadrados por cada temporada de crecimiento. Cuando se aplica como se indica, este producto provee 0.75 libras de nitrógeno por cada 1000 pies cuadrados. Comuníquese con su Servicio de Extensión Cooperativa local para obtener información específica para su región geográfica.

CUANDO APLICARLO: Aplíquele cuando las malezas de hojas anchas estén creciendo activamente en la primavera o comienzo del otoño. Aplíquelo cuando el viento esté calmado (ojo un día tranquilo) para asegurar una cobertura uniforme y evitar que los gránulos se esparzan hacia flores, vegetales y adornos ornamentales. Es posible que sea necesario repetir el tratamiento si el césped está invadido por las malezas, si prevalecen las condiciones adversas de poca humedad o si las malezas están en un estado de poco crecimiento.

INSTRUCCIONES PARA LA PLANCHA:

1. Pode el césped a una altura normal 1 o 2 días antes de la aplicación.
2. Riegue el césped ahundantemente al menos 1 o 2 días antes de la aplicación para mantener la humedad del suelo hasta la próxima sesión de riego (vea el paso 5).
3. Aplíquelo cuando las malezas de hojas anchas estén jóvenes y creciendo activamente, preferiblemente en la mañana cuando el rocío esté aún sobre la hierba. En el momento de la aplicación, la humedad del rocío, la lluvia o el riego sobre las hojas de la maleza ayudará al control. Si la hierba no está húmeda al momento de la aplicación, rocíela ligeramente con agua antes de aplicar el producto.
4. Evite aplicarlo cuando las temperaturas no sean consistentemente superiores a los 90°F (32°C).
5. No permita que el agua entre el producto de las hojas de la maleza por 1 o 2 días después de la aplicación. Sólo entonces, se debe aplicar una sesión de riego abundante.

AJUSTES DE LOS ESPARCIDORES

Aplíquelo siempre con un esparcidor giratorio o por goteo para distribuir los gránulos uniformemente. No lo aplique nunca a mano.

IMPORTANTE: Es muy importante que cumpla con las indicaciones que se incluyen con este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede ocasionar daños al césped. Los ajustes del esparcidor son aproximados y puede haber diferencias debido a las condiciones del esparcidor, la velocidad con la que usted lo aplica y las irregularidades del área tratada. Asegúrese de revisar el esparcidor antes de comenzar y ajústalo si fuera necesario. Consulte el manual del propietario. Nunca lo aplique a mano.

Tamaño de la bolsa (lbs)	Tasa de aplicación (lbs / 1,000 sq. ft.)	Area tratada por una bolsa de este tamaño (sq. ft.)
12.5	2.5	5,000

DISTRIBUCIÓN CON UN SOLO PASE:

- Para esparcidores giratorios: Trabaje en intervalos correspondientes con el ancho de la franja de tierra.
- Para esparcidores por goteo: Solape la trayectoria de las rueda para garantizar una cobertura completa.
- Para lograr mejores resultados, aplique el producto uniformemente y trate de no dejar franjas sin cubrir, saltarse espacios o aplicar dos veces en una misma área. Camine con un paso estable para asegurar una cobertura uniforme. Cierre la abertura del esparcidor cuando gire para cambiar de dirección o cuando se detenga.

INSTRUCCIONES PARA LA CALIBRACIÓN Y AJUSTES DEL ESPARCIDOR: Calibre el esparcidor según las instrucciones del fabricante. El ajuste inicial del esparcidor pudiera requerir un reglado para aplicar la tasa recomendada a partir de las condiciones reales de aplicación. Para garantizar una distribución apropiada, calibre su esparcidor antes de la aplicación.

Para 2.5 libras de producto por 1000 pies cuadrados de césped: Verifique su esparcidor en 400 pies cuadrados y ajústelo para aplicar a una tasa de 1 libra de producto por 400 pies cuadrados.

ESPARCIDORES POR DISPERSIÓN: Para lograr mejores resultados, aplíquelo en una trayectoria circular comenzando por la parte exterior y trabajando hacia el centro, solapando ligeramente sobre el área recién aplicada para garantizar una cobertura completa.

ESPARCIDORES POR GOTEO: Aplíquelo en 2 franjas a lo largo de cada extremo y aplíquelo luego a lo largo del césped, solapando ligeramente sobre el área recién aplicada y cerrando el esparcidor antes de cada giro o al final de cada trayectoria por los bordes.

AJUSTE DEL ESPARCIDOR PARA CÉSPED

ESTOS AJUSTES SON SOLO UNA GUÍA. Fueron todos determinados por profesores profesionales usando esparcidores nuevos. Las condiciones de su esparcidor, la velocidad con la que usted camina, la ligereza de su césped o incluso la humedad pueden afectar la velocidad con la que sale el nutriente de su esparcidor. Preste atención al flujo; si el fertilizante sale más rápido o más despacio de lo que usted cubre el césped, modifique el ajuste del esparcidor. Cierre siempre el esparcidor cuando lo rellene, cuando gire o cuando se detenga.

Esparcidor	Ajuste para aplicar 2.5 lbs / 1000 sq. ft.
Esparcidores por goteo Scotts®	3.5
Esparcidores por giratorio Scotts®	3.5
Esparcidores por dispersión Sta-Green®	3.25
Esparcidores por dispersión Pennington®	3

ALMACENAJE Y DESECHO

No contamine el agua, los alimentos ni el pienso al almacenar y desechar el producto.

ALMACENAJE DE PESTICIDAS: Guárdelo en el recipiente original y mantenga las bolsas que contienen el producto sin usar en un área de almacenaje cerrada e inaccesible para niños y animales domésticos.

DESECHO DE PESTICIDAS Y MANIPULACIÓN DEL RECIPIENTE: Este recipiente no es rellenable. No lo reutilice ni lo rellene. Si está vacío: Tírelo a la basura o recíclelo si el servicio está disponible. Si está parcialmente lleno: Llame a su agencia local para el desecho de sustancias sólidas para recibir instrucciones en cuanto al desecho. No vierta nunca el producto sin usar en vertidos de desagüe interiores o exteriores.

GARANTÍA LIMITADA Y RENUNCIA DE RESPONSABILIDAD

GARANTÍA Y RENUNCIA DE RESPONSABILIDAD: En consonancia con lo estipulado por la ley correspondiente, Central Garden & Pet y la División de Jardinería garantizan este producto de calidad hasta la cantidad del precio de compra y solo cuando se usa como se indica.

30-0-4

ANÁLISIS GARANTIZADO

Total de nitrógeno (N)	30%
10% de nitrógeno urea*	
Potasa soluble (K2O)	4%
Calcio (Ca)	1%

*Derivado de la urea, una con revestimiento de polímero, mezclas de potasa y cal.

*Contiene un 12% de nitrógeno urea de liberación lenta a partir de la urea con revestimiento de polímero.

*Contiene un 18% de nitrógeno estabilizado con tiosulfato N-(n-butil) de triamida.

**El nitrógeno estabilizado reduce la pérdida de nitrógeno hacia el ambiente y mejora su eficiencia.

CONTIENE TAMBIEN INGREDIENTES QUE NO SON NUTRIENTES DE PLANTAS

ANÁLISIS GARANTIZADO PARA ENMIENDAS DE SUELO

Ingredientes activos:

Hongos micorrizas	1.12 unos por
Glomus intraradices	0.28 unos por
Glomus mosseae	0.28 unos por
Glomus aculeatum	0.28 unos por
Glomus aggregatum	0.28 unos por

Ingredientes inertes (como enmienda del suelo):

98.34% de ingredientes inertes

1.184% de ingredientes herbicidas

Para ver la información disponible acerca del contenido y los niveles de metales en este producto, visite la página de Internet: http://www.aapfco.org/metals.htm

Ciertos estados tienen límites en cuanto a las tasas y periodos de aplicación de fertilizantes. Consulte con su Servicio de Extensión Cooperativa local para conocer la información actualizada sobre los límites anuales de aplicación de fertilizantes de césped en su región.

La información acerca de los materiales primas que componen este producto se puede obtener escribiendo al vendedor mencionado en este etiquetado y refiriendo el número de lote que se encuentra en este recipiente.

AVISO: Este producto contiene sustancias quimicas reconocidas en el estado de California como causantes de cáncer y defectos congénitos u otros daños al sistema reproductivo.

† **Garantía de satisfacción de Pennington:** Si, después de haber seguido cuidadosamente las instrucciones, usted no está satisfecho con los resultados del producto, envíe su nombre, dirección y prueba de la compra (recibo original con fecha) a:

Pennington Brand
1000 Parkwood Circle, Suite 700, Atlanta, GA 30339

Pennington and Pennington with design are registered trademarks of Pennington Seed, Inc. Central Garden & Pet, Central Garden & Pet with design and UltraGreen are trademarks of Central Garden & Pet Company. Scotts, Turf Builder, and Edgeguard are registered trademarks of ORO Investments, Inc. Sta-Green is a registered trademark of LLC. ©2015 Central Garden & Pet Company.

Guaranteed & Distributed by Guaranteed per Central Garden & Pet Garden Division, 1000 Parkwood Circle, Suite 700, Atlanta, GA 30339

gardendivisioninfo@centralgarden.com
www.penningtonfertilizer.com
1-800-205-5761

EPA Reg. No. 2217-987-10008
EPA Est. No. 35614-GA-1, 35817-OR-1, 33783-OH-1, 37263-CO-1
Solicitud usted es the First letter of Rule Code or Container

0 21496 01327 5

CENTRAL Garden & Pet

3005170148 -07/15 • UG80 v.14

Expiration Date

This herbicide controls weeds listed in the following table / Este herbicida controla las malezas mencionadas en la siguiente tabla:

Acacia(fr. spurge)	Burweed, lawn	Cocklebur, oriental	Fleabane, rough	Pusley, large flower	Spurge, spotted	Virginia creeper
Acacia(fr. slender	(spineweed)	Cocklebur, spiny	Galinsoga, hairy	Ragweed, lac	Starwort	Welcome cassia
Acacia, sicklepod	Buttercup, bulbous	Cocklebur, tropical	Galinsoga, smallflower	Ragweed, common	Stinging nettle	Wild carrot
Acder, beach	Buttercup, common	Cocklebur, common	Geranium,	Ragweed, lacelaf	Strawberry, India mock	Wild cucumber
Acter, purple	Buttercup, creeping	Combseed,	Geranium, cut-leaf	Ragweed, western	Sump rayless	Wild lettuce
Acter, white heath	(spotted)	Creeping Bentgrass	Geranium,	Ragweed, ragweed	Tansy	Wild oat
Acter, white prairie	Buttercup, virgina	Creeping Bentgrass	Germander,	Rocket, sweet	Thistle, blessed	Wild four-o-clock
Acter, white wood	Cabbage,	(Creeping charlie)	Gumweed	Rough cocklebur	Thistle, bull	Wild garlic
Dandelion, Catchweed	Carpetweed	Crabgrass	Hawkweed, meadow	Rough fleabane	Thistle, canada	Wild lettuce
Dandelion, Northern	Carpetweed	Crabgrass, smooth	Hawkweed, meadow	Sheepsorrel	Thistle, field	Wild mustard
Dandelion, smooth	Catsip	Crabgrass,	Hawkweed, orange	Smartweed, pink	Thistle, flodman	Wild oat
Dandelion, smooth	Catsear, spotted	Dandelion, common	Hawkweed, yellow	Smartweed, pennsylvania	Thistle, musk	Wild radish
Dock, curly	Chamber bitter	Dayflower,	Henbit	Smartweed,	Thistle, plumless	Wild violet
Dollar, Florida	Chamber bitter	Dayflower, common	Horsetail dryness	Smartweed,	Thistle, plumeless	Willowherb
Endweed, field	Chickweed, common	Dayflower,	Horsetail	Sorrel, red	Thistle, russian	Wild radish
Endweed, hedge	Chickweed, mouseear	Dichondra	Horsetail	Sorrel, red	Thistle, scotch	Wild rape
Etter Wintercress	Chickweed, sticky	Dock, broadleaf	Hours cress	Sorrel,	Thistle, waystalf	Wild strawberry
Etter Wintercress	Clover,	Dock, curly	Hours cress	Speedwell,	Thistle, wild	Wild violet
Eliflamerus	Clover, alsike	Dogfennel	Horsweed	Speedwell,	Thistlebur	Wild violet
Black Medic	Clover, alsike	Dogfennel	Jimsonweed	Speedwell, bi-lobed	Trailing crownvetch	Wild violet
Black-eyed Susan	Clover, crimson	Dollar weed	Jimsonweed	Speedwell, common	Velvetleaf	Willowherb, creeping
Blast flower redweed	Clover, hop	(Pennywort)	Knawel	Speedwell, corn	Venus, blue	Willowherb, yellow
Bracts, red	Clover, hop	Dollar weed	Knotweed, prostrate	Speedwell, corn	Venus, perfoliate	Woodsorrel
Bracts,yellow	Clover, red	Dollarweed	Knotweed,	Speedwell, persian	Venus, prophylate	Woody croton
Briefly sabrgrass	Clover, strawberry	Dollarweed	Lambsquarters	Speedwell, purslane	Venus, tall	Wormseed
Broomweed	Clover, white	Elephantopus	Lamiastrum	Speedwell, slender	Vetch, bird	Yarrow, common
Buckhorn	Clover, white sweet	Elephantopus	Mallow, alcea	Speedwell, snow	Vetch, hairy	Yarrow, western
Buckhorn	Condo, yellow sweet	Fallow, wild	Mallow, common	Speedwell, speedwell	Vetch,	Yarrow, yellow
Burdock	Condo, white	Fallow, common	Mallow, cornfield	Speedwell, speedwell	Vetch, narrowleaf	
Burdock	Cocklebur	Fleabane, daisy	Mallow, dwarf	Spurge, prostrate	Vetch, wild	
		(annual)				

			Pepperweed, virginia	Pusley, large flower	Spurge, spotted	Virginia creeper	
			Marestail	Ragweed, lac	Starwort	Welcome cassia	
			Mallow, venice	Ragweed, common	Stinging nettle	Wild carrot	
			Mallowweed	Ragweed, lacelaf	Strawberry, India mock	Wild cucumber	
			Mallowweed	Ragweed, western	Sump rayleaf	Wild lettuce	
			Menagerie, bigseed	Ragweed, ragweed	Tansy	Wild oat	
			Menagerie, bigseed	Rocket, sweet	Thistle, blessed	Wild four-o-clock	
			Menagerie, bigseed	Rough cocklebur	Thistle, bull	Wild garlic	
			Menagerie, bigseed	Rough fleabane	Thistle, canada	Wild lettuce	
			Menagerie, bigseed	Sheepsorrel	Thistle, field	Wild mustard	
			Menagerie, bigseed	Smartweed, pink	Thistle, flodman	Wild oat	
			Menagerie, bigseed	Smartweed,	Thistle,	Wild radish	
			Menagerie, bigseed	Smartweed, pennsylvania	Thistle, musk	Wild violet	
			Menagerie, bigseed	Sorrel, red	Thistle, plumless	Willowherb	
			Menagerie, bigseed	Sorrel, red	Thistle, plumeless	Wild radish	
			Menagerie, bigseed	Sorrel, red	Thistle, russian	Wild rape	
			Menagerie, bigseed	Speedwell,	Thistle, scotch	Wild strawberry	
			Menagerie, bigseed	Speedwell,	Thistle, waystalf	Wild violet	
			Menagerie, bigseed	Speedwell,	Thistle, wild	Wild violet	
			Menagerie, bigseed	Speedwell, bi-lobed	Thistlebur	Willowherb,	
			Menagerie, bigseed	Speedwell, common	Trailing crownvetch	Willowherb, creeping	
			Menagerie, bigseed	Speedwell, corn	Velvetleaf	Willowherb, yellow	
			Menagerie, bigseed	Speedwell, corn	Venus, blue	Woodsorrel,	
			Menagerie, bigseed	Speedwell, persian	Venus, perfoliate	Woody croton	
			Menagerie, bigseed	Speedwell, purslane	Venus, prophylate	Wormseed	
			Menagerie, bigseed	Speedwell, slender	Venus, tall	Yarrow, common	
			Menagerie, bigseed	Speedwell, snow	Vetch, hairy	Yarrow, western	
			Menagerie, bigseed	Speedwell, speedwell	Vetch,	Yarrow, yellow	
			Menagerie, bigseed	Spurge, prostrate	Vetch, narrowleaf		
			Menagerie, bigseed	Spurge, prostrate	Vetch, wild		